



Nonlinear Donati compatibility conditions for the nonlinear Kirchhoff -von K arm an-Love plate theory

Philippe G. Ciarlet, Giuseppe Geymonat, Francoise Krasucki

► To cite this version:

Philippe G. Ciarlet, Giuseppe Geymonat, Francoise Krasucki. Nonlinear Donati compatibility conditions for the nonlinear Kirchhoff -von K arm an-Love plate theory. Comptes rendus hebdomadaires des séances de l'Académie des sciences, 2013, 351, pp.405-409. 10.1016/j.crma.2013.05.012 . hal-00959792

HAL Id: hal-00959792

<https://hal.science/hal-00959792>

Submitted on 16 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nonlinear Donati compatibility conditions for the nonlinear Kirchhoff-von Kármán-Love plate theory

Philippe G. Ciarlet^a, Giuseppe Geymonat^b, Francoise Krasucki^c

^a*Department of Mathematics, City University of Hong Kong, 83 Tat Chee Avenue, Kowloon, Hong Kong*

^b*Laboratoire de Mécanique des Solides, UMR 7649-0176, Ecole Polytechnique, 91128 Palaiseau Cedex, France*

^c*I3M, UMR-CNRS 5149, Université de Montpellier II, Place Eugène Bataillon, 34095 Montpellier Cedex 5, France*

Received *****

Presented by Philippe G. Ciarlet

Abstract

Let ω be a simply-connected domain in \mathbb{R}^2 and let $(E_{\alpha\beta})$ and $(F_{\alpha\beta})$ be two symmetric 2×2 matrix fields with components in $L^2(\omega)$. In this Note, we identify nonlinear compatibility conditions “of Donati type” that the components $E_{\alpha\beta}$ and $F_{\alpha\beta}$ must satisfy in order that there exists a vector field $(\eta_1, \eta_2, w) \in H_0^1(\omega) \times H_0^1(\omega) \times H_0^2(\omega)$ such that

$$\frac{1}{2}(\partial_\alpha \eta_\beta + \partial_\beta \eta_\alpha + \partial_\alpha w \partial_\beta w) = E_{\alpha\beta} \text{ and } \partial_{\alpha\beta} w = F_{\alpha\beta} \text{ in } \omega.$$

The left-hand sides of these relations are the components of tensors found in the Kirchhoff-von Kármán-Love theory of nonlinearly elastic plates.

Résumé

Conditions de compatibilité non linéaires de Donati pour la théorie non linéaire des plaques de Kirchhoff-von Kármán-Love. Soit ω un domaine simplement connexe de \mathbb{R}^2 et soit $(E_{\alpha\beta})$ et $(F_{\alpha\beta})$ deux champs de matrices 2×2 symétriques dont les composantes sont dans $L^2(\omega)$. Dans cette Note, on identifie et justifie des conditions non linéaires de compatibilité “de type Donati” que doivent satisfaire les composantes $E_{\alpha\beta}$ et $F_{\alpha\beta}$ afin qu’il existe un champs de vecteurs $(\eta_1, \eta_2, w) \in H_0^1(\omega) \times H_0^1(\omega) \times H_0^2(\omega)$ tel que

$$\frac{1}{2}(\partial_\alpha \eta_\beta + \partial_\beta \eta_\alpha + \partial_\alpha w \partial_\beta w) = E_{\alpha\beta} \text{ et } \partial_{\alpha\beta} w = F_{\alpha\beta} \text{ dans } \omega.$$

Les membres de gauche de ces relations sont les composantes de tenseurs trouvés dans la théorie de Kirchhoff-von Kármán-Love des plaques non linéairement élastiques.

Email addresses: mapgc@cityu.edu.hk (Philippe G. Ciarlet), giuseppe.geymonat@lms.polytechnique.fr (Giuseppe Geymonat), krasucki@math.univ-montp2.fr (Francoise Krasucki).

1. Preliminaries

Greek indices vary in $\{1, 2\}$ and the convention summation with respect to repeated indices is used. A *domain* in \mathbb{R}^2 is a bounded, open, and connected subset ω of \mathbb{R}^2 with a Lipschitz-continuous boundary $\partial\omega$, the set ω being locally on the same side of $\partial\omega$. Partial derivatives of the first, second, and third, order of functions of $y = (y_\alpha) \in \omega$ are denoted $\partial_\alpha := \partial/\partial y_\alpha$, $\partial_{\alpha\beta} := \partial^2/\partial y_\alpha \partial y_\beta$, and $\partial_{\alpha\beta\sigma} := \partial^3/\partial y_\alpha \partial y_\beta \partial y_\sigma$; the same notations are used for partial derivatives in the sense of distributions.

Vector fields and matrix fields, and spaces of vector fields, defined over ω are denoted by boldface letters. The usual Sobolev spaces over ω are denoted $H^m(\omega)$, $m \in \mathbb{Z}$, and $H_0^m(\omega)$, $m \geq 1$; the norm in $H^m(\omega)$, $m \in \mathbb{Z}$, is denoted $\|\cdot\|_{m,\omega}$; in particular then, $\|\cdot\|_{0,\omega}$ is the norm of $H^0(\omega) = L^2(\omega)$. The notation $\mathbb{L}^2(\omega)$ designates the space of all 2×2 *symmetric* matrix fields with components in $L^2(\omega)$. If $\mathbf{S} = (S_{\alpha\beta})$ is a 2×2 matrix field with smooth enough components defined over ω , its *divergence* $\mathbf{div} \mathbf{S}$ is the vector field defined by $(\mathbf{div} \mathbf{S})_\alpha = \partial_\alpha S_{\alpha\beta}$.

If X and Y are two (real) vector spaces and A is a linear operator from X to Y ,

$$\text{Im } A := \{y \in Y; y = Ax \text{ for at least one } x \in X\} \text{ and } \text{Ker } A := \{x \in X; Ax = 0\}.$$

The notation $X' \langle \cdot, \cdot \rangle_X$ designates the duality between a normed vector space X and its dual X' .

In the classical *Kirchhoff-von Kármán-Love theory of nonlinearly elastic plates* (see, e.g., Chapters 4 and 5 in [2]), the unknown displacement field of the middle surfac $\bar{\omega}$ of the plate minimizes an energy whose integrand contains a positive-definite quadratic function of the *change of metric* and *change of curvature tensors*, respectively defined by

$$E_{\alpha\beta} := \frac{1}{2}(\partial_\alpha \eta_\beta + \partial_\beta \eta_\alpha + \partial_\alpha w \partial_\beta w) \text{ and } F_{\alpha\beta} := \partial_{\alpha\beta} w, \quad (1)$$

for an arbitrary displacement field $(\eta_1, \eta_2, w) \in H_0^1(\omega) \times H_0^1(\omega) \times H_0^2(\omega)$ (we consider here plates that are clamped over their entire lateral face).

In the *intrinsic* approach to the same theory, the matrix fields $(E_{\alpha\beta}) \in \mathbb{L}^2(\omega)$ and $(F_{\alpha\beta}) \in \mathbb{L}^2(\omega)$ are considered as the *sole unknowns*. There thus arises the question as to whether there exist suitable *compatibility conditions* that the components $E_{\alpha\beta}$ and $F_{\alpha\beta}$ of these matrix fields should satisfy in order that there exists a vector field $(\eta_1, \eta_2, w) \in H_0^1(\omega) \times H_0^1(\omega) \times H_0^2(\omega)$ satisfying (1). As shown in [6], if the domain ω is simply-connected, the **nonlinear Saint-Venant compatibility conditions**

$$\begin{aligned} \partial_{\sigma\tau} E_{\alpha\beta} + \partial_{\alpha\beta} E_{\sigma\tau} - \partial_{\alpha\sigma} E_{\beta\tau} - \partial_{\beta\tau} E_{\alpha\sigma} + F_{\alpha\beta} F_{\sigma\tau} - F_{\alpha\sigma} F_{\beta\tau} &= 0 \text{ in } H^{-2}(\omega), \\ \partial_\sigma F_{\alpha\beta} - \partial_\beta F_{\alpha\sigma} &= 0 \text{ in } H^{-1}(\omega), \end{aligned}$$

constitute one possible answer to this question. The objective of this Note is to give (cf. Theorem 4.2) a different answer to the same question, this time in the form of *variational equations*, which as such constitute examples of **nonlinear Donati compatibility conditions** (a general presentation of Saint-Venant and Donati compatibility conditions as they arise in three-dimensional linearized elasticity is found in Chapter 6 in [3]).

Complete proofs and an application to intrinsic nonlinear plate theory will be found in [5].

2. An existence theorem for an Airy-function

The following result is a “weak” version (already used in [6]) of a classical result for smooth functions. Its proof is based on the two-dimensional version of the *weak Poincaré lemma* due to [4] and then given a substantially simpler proof in [10]; cf. also Thm. 6.17-4 in [3].

Theorem 2.1 *Let ω be a simply-connected domain in \mathbb{R}^2 , and let there be given a matrix field $(F_{\alpha\beta}) \in \mathbb{L}^2(\omega)$ that satisfies*

$$\partial_\sigma F_{\alpha\beta} - \partial_\beta F_{\alpha\sigma} = 0 \text{ in } H^{-1}(\omega).$$

Then there exists a function $\varphi \in H^2(\omega)$, unique up to the addition of a polynomial of degree ≤ 1 , such that

$$\partial_{\alpha\beta}\varphi = F_{\alpha\beta} \text{ in } L^2(\omega).$$

Theorem 2.1 can be immediately recast as an existence result of an ad hoc *Airy function* (denoted φ in the next theorem) under low regularity assumptions. As such, it complements Thm. 2 of [7], where the existence of an Airy function was established, again in the space $H^2(\omega)$, but for *non-simply-connected domains*, under the assumption that the tensor field noted \mathbf{S} in the next theorem satisfies in addition the usual global equilibrium equations.

Theorem 2.2 *Let ω be a simply-connected domain in \mathbb{R}^2 , and let there be given a matrix field $\mathbf{S} = (S_{\alpha\beta}) \in \mathbb{L}^2(\omega)$ that satisfies*

$$\operatorname{div} \mathbf{S} = \mathbf{0} \text{ in } H^{-1}(\omega).$$

Then there exists a function $\varphi \in H^2(\omega)$, unique up to the addition of a polynomial of degree ≤ 1 , such that

$$\partial_{11}\varphi = S_{22}, \quad \partial_{12}\varphi = -S_{12}, \quad \partial_{22}\varphi = S_{11} \text{ in } L^2(\omega). \quad (2)$$

A function φ satisfying the relations (2) is called an **Airy function associated with the matrix field \mathbf{S}** .

3. Linear Donati compatibility conditions for linearly elastic plates

For convenience, we consider separately the existence of the “horizontal” components η_α , and that of the “vertical” component w of the unknown vector field.

Theorem 3.1 *Let ω be a domain in \mathbb{R}^2 and let there be given a matrix field $(e_{\alpha\beta}) \in \mathbb{L}^2(\omega)$ that satisfies*

$$\int_\omega e_{\alpha\beta} s_{\alpha\beta} \, dy = 0 \text{ for all } (s_{\alpha\beta}) \in \mathbb{L}^2(\omega) \text{ such that } \partial_\alpha s_{\alpha\beta} = 0 \text{ in } H^{-1}(\omega). \quad (3)$$

Then there exists a vector field $(\eta_\alpha) \in H_0^1(\omega) \times H_0^1(\omega)$ such that

$$\frac{1}{2}(\partial_\alpha \eta_\beta + \partial_\beta \eta_\alpha) = e_{\alpha\beta} \text{ in } L^2(\omega),$$

and such a vector field (η_α) is uniquely determined.

Proof. See, e.g., [1], or [8] and [9]. □

Theorem 3.2 *Let ω be a domain in \mathbb{R}^2 and let there be given a matrix field $(F_{\alpha\beta}) \in \mathbb{L}^2(\omega)$ that satisfies*

$$\int_\omega F_{\alpha\beta} T_{\alpha\beta} \, dy = 0 \text{ for all } (T_{\alpha\beta}) \in \mathbb{L}^2 \text{ such that } \partial_{\alpha\beta} T_{\alpha\beta} = 0 \text{ in } H^{-2}(\omega).$$

Then there exists a function $w \in H_0^2(\omega)$ such that

$$\partial_{\alpha\beta} w = F_{\alpha\beta} \text{ in } L^2(\omega), \quad (4)$$

and this function is uniquely determined.

Sketch of proof. Let the continuous linear operator $\mathbf{H} : H_0^2(\omega) \rightarrow \mathbb{L}^2(\omega)$ be defined by

$$\mathbf{H}w = \begin{pmatrix} \partial_{11}w & \partial_{12}w \\ \partial_{21}w & \partial_{22}w \end{pmatrix} \in \mathbb{L}^2(\omega) \text{ for each } w \in H_0^2(\omega).$$

Then one shows that $\text{Im } \mathbf{H}$ is a closed subspace of $\mathbb{L}^2(\omega)$, and that the dual operator of \mathbf{H} is $\text{div } \mathbf{div} : \mathbb{L}^2(\omega) \rightarrow H^{-2}(\omega)$. The conclusion then follows from Banach closed range theorem. \square

Relation (3) and (4) constitute the **linear Donati compatibility conditions** corresponding to the Kirchhoff-Love theory of linearly elastic plates. Note that they hold regardless of whether the domain ω is simply-connected.

4. Nonlinear Donati compatibility conditions for nonlinearly elastic plates

The Green's formula (5) established in the next theorem is crucial to our subsequent analysis.

Theorem 4.1 For all functions $w \in H_0^2(\omega)$ and $\varphi \in H^2(\omega)$,

$$\int_{\omega} (\partial_{11}w \partial_{22}w - \partial_{12}w \partial_{12}w) \varphi \, dy = \int_{\omega} \left\{ -\frac{1}{2}(\partial_1 w)^2 \partial_{22}\varphi - \frac{1}{2}(\partial_2 w)^2 \partial_{11}\varphi + \partial_1 w \partial_2 w \partial_{12}\varphi \right\} dy. \quad (5)$$

Sketch of proof. Both sides of (5) being continuous functions of $(w, \varphi) \in H_0^2(\omega) \times H^2(\omega)$, it is enough to establish (5) for all $(w, \varphi) \in \mathcal{D}(\omega) \times H^2(\omega)$. To this end, one uses the integration by parts formulas in Sobolev spaces. \square

The next theorem constitutes the main result of this Note.

Theorem 4.2 Let ω be a simply-connected domain in \mathbb{R}^2 . Given a matrix field $\mathbf{S} \in \mathbb{L}^2(\omega)$ that satisfies $\text{div } \mathbf{S} = \mathbf{0}$ in $\mathbf{H}^{-1}(\omega)$, there exists a unique function $\varphi \in H^2(\omega)$ such that (cf. Theorem 2.2)

$$\partial_{11}\varphi = S_{22}, \quad \partial_{12}\varphi = -S_{12}, \quad \partial_{22}\varphi = S_{11} \text{ in } L^2(\omega) \text{ and } \int_{\omega} \varphi \, dy = \int_{\omega} \partial_{\alpha}\varphi \, dy = 0.$$

Let

$$\Phi : \{\mathbf{S} \in \mathbb{L}^2(\omega); \text{div } \mathbf{S} = \mathbf{0} \text{ in } \mathbf{H}^{-1}(\omega)\} \rightarrow \{\psi \in H^2(\omega); \int_{\omega} \psi \, dy = \int_{\omega} \partial_{\alpha}\psi \, dy = 0\}$$

denote the mapping defined in this fashion, i.e., by $\Phi(\mathbf{S}) := \varphi$.

Let there be given two matrix fields $(E_{\alpha\beta}) \in \mathbb{L}^2(\omega)$ and $\mathbf{F} = (F_{\alpha\beta}) \in \mathbb{L}^2(\omega)$ that satisfy

$$\int_{\omega} F_{\alpha\beta} T_{\alpha\beta} \, dy = 0 \text{ for all } \mathbf{T} = (T_{\alpha\beta}) \in \mathbb{L}^2(\omega) \text{ such that } \text{div } \mathbf{div } \mathbf{T} = \mathbf{0} \text{ in } H^{-2}(\omega), \quad (6)$$

$$\int_{\omega} \{E_{\alpha\beta} S_{\alpha\beta} + (\det \mathbf{F}) \Phi(\mathbf{S})\} \, dy = 0 \text{ for all } \mathbf{S} = (S_{\alpha\beta}) \in \mathbb{L}^2(\omega) \text{ such that } \text{div } \mathbf{S} = \mathbf{0} \text{ in } \mathbf{H}^{-1}(\omega). \quad (7)$$

Then there exists a vector field $(\eta_1, \eta_2, w) \in H_0^1(\omega) \times H_0^1(\omega) \times H_0^2(\omega)$ such that

$$\begin{aligned} \frac{1}{2}(\partial_{\alpha}\eta_{\beta} + \partial_{\beta}\eta_{\alpha} + \partial_{\alpha}w \partial_{\beta}w) &= E_{\alpha\beta} \text{ in } L^2(\omega), \\ \partial_{\alpha\beta}w &= F_{\alpha\beta} \text{ in } L^2(\omega), \end{aligned}$$

and such a vector field (η_1, η_2, w) is uniquely determined.

Proof. Relation (6) shows that there exists a uniquely determined function $w \in H_0^2(\omega)$ such that (cf. Theorem 3.2)

$$F_{\alpha\beta} = \partial_{\alpha\beta}w \text{ in } L^2(\omega). \quad (8)$$

Given any matrix field $\mathbf{S} = (S_{\alpha\beta}) \in \mathbb{L}^2(\omega)$ such that $\mathbf{div} \mathbf{S} = \mathbf{0}$ in $\mathbf{H}^{-1}(\omega)$, there exists a uniquely determined function $\varphi \in H^2(\omega)$ such that (cf. Theorem 2.2)

$$S_{11} = \partial_{22}\varphi, \quad S_{12} = -\partial_{12}\varphi, \quad S_{22} = \partial_{11}\varphi \text{ in } L^2(\omega).$$

Therefore, for any such matrix field \mathbf{S} ,

$$\int_{\omega} E_{\alpha\beta} S_{\alpha\beta} \, dy = \int_{\omega} \{E_{11}\partial_{22}\varphi + E_{22}\partial_{11}\varphi - 2E_{12}\partial_{12}\varphi\} \, dy,$$

and (cf. (7))

$$\begin{aligned} \int_{\omega} (\det \mathbf{F}) \Phi(\mathbf{S}) \, dy &= \int_{\omega} (F_{11}F_{22} - (F_{12})^2) \varphi \, dy, \\ &= \int_{\omega} \{(\partial_{11}w\partial_{22}w - \partial_{12}w\partial_{12}w) \varphi \, dy\}. \end{aligned}$$

The left-hand side of relation (6) can thus be rewritten as (cf. 5)

$$\begin{aligned} &\int_{\omega} \{E_{\alpha\beta} S_{\alpha\beta} + (\det \mathbf{F}) \Phi(\mathbf{S})\} \, dy \\ &= \int_{\omega} \left\{ \left(E_{11} - \frac{1}{2}(\partial_1 w)^2 \right) \partial_{22}\varphi - 2 \left(E_{12} - \frac{1}{2}\partial_1 w \partial_2 w \right) \partial_{12}\varphi + \left(E_{22} - \frac{1}{2}(\partial_2 w)^2 \right) \partial_{11}\varphi \right\} \, dy \\ &= \int_{\omega} \left\{ \left(E_{11} - \frac{1}{2}(\partial_1 w)^2 \right) S_{11} + 2 \left(E_{22} - \frac{1}{2}\partial_1 w \partial_2 w \right) S_{12} + \left(E_{22} - \frac{1}{2}(\partial_2 w)^2 \right) S_{22} \right\} \, dy. \end{aligned}$$

Since this last relation holds for all $\mathbf{S} = (S_{\alpha\beta}) \in \mathbb{L}^2(\omega)$ such that $\mathbf{div} \mathbf{S} = \mathbf{0}$ in $\mathbf{H}^{-1}(\omega)$, there exists a uniquely determined vector field $(\eta_{\alpha}) \in H_0^1(\omega) \times H_0^1(\omega)$ such that (cf. Theorem 3.1)

$$E_{\alpha\beta} - \frac{1}{2}\partial_{\alpha}w\partial_{\beta}w = \frac{1}{2}(\partial_{\alpha}\eta_{\beta} + \partial_{\beta}\eta_{\alpha}) \text{ in } L^2(\omega).$$

This completes the proof. \square

Relations (6) and (7) constitute the **nonlinear Donati compatibility conditions** corresponding to the Kirchhoff-von Kármán-Love theory of nonlinearly elastic plates. Note that, when properly extended, they can also cover the case where the domain ω is not simply-connected; cf. [5].

Finally, note that, as expected, the linearization of the nonlinear Donati compatibility conditions (7) reduce to the linear ones (cf. (3)). \square

Acknowledgment

This work was partially supported by a grant from the Research Grants Council of the Hong Kong Special Administrative Region, China [Project No. 9041738- CityU 100612].

References

- [1] C. Amrouche, P.G. Ciarlet, L. Gratie, S. Kesavan, On the characterizations of matrix fields as linearized strain tensor fields, J. Math. Pures Appl. 86 (2006) 116–132.
- [2] P.G. Ciarlet, Mathematical Elasticity, Volume II: Theory of Plates, North-Holland, Amsterdam 1997.
- [3] P.G. Ciarlet, Linear and Nonlinear Functional Analysis with Applications, SIAM, 2013.

- [4] P.G. Ciarlet, P. Ciarlet, Jr., Another approach to linearized elasticity and a new proof of Korn's inequality, *Math. Models Methods Appl. Sci.* 15 (2005) 259–271.
- [5] P.G. Ciarlet, G. Geymonat, F. Krasucki, Nonlinear Donati compatibility conditions and the intrinsic approach for nonlinearly elastic plates, in preparation.
- [6] P.G. Ciarlet, S. Mardare, Nonlinear Saint-Venant compatibility conditions and the intrinsic approach for nonlinearly elastic plates, *Math. Models Methods Appl. Sci.*, to appear.
- [7] G. Geymonat, F. Krasucki, On the existence of the Airy function in Lipschitz domains. Application to the traces of H^2 , *C.R. Acad. Sci. Paris, Sér. I* 330 (2000) 355–360.
- [8] G. Geymonat, F. Krasucki, Some remarks on the compatibility conditions in elasticity, *Accad. Naz. Sci. XL* 123 (2005) 175–182.
- [9] G. Geymonat, F. Krasucki, Hodge decomposition for symmetric matrix fields and the elasticity complex in Lipschitz domains, *Comm. Pure Appl. Anal.* 8 (2009) 295–309.
- [10] S. Kesavan, On Poincaré's and J.L. Lions' lemmas, *C.R. Acad. Sci. Paris, Sér. I* 340 (2005) 27–30.